arXiv:1602.06288v2 [math.CA] 11 Oct 2016

Some Approximation Results by (p,q)-analogue of Bernstein-Stancu Operators (Revised)

M. Mursaleen, **Khursheed J. Ansari** and **Asif Khan**
Department of Mathematics, Aligarh Muslim University, Aligarh–202002, India
mursaleenm@gmail.com; ansari.jkhursheed@gmail.com; asifjnu07@gmail.com

Abstract

In this paper, We have given a corrigendum to our paper "Some Approximation Results by (p,q)-analogue of Bernstein-Stancu Operators" published in Applied Mathematics and Computation 264 (2015) 392-402. We introduce a new analogue of Bernstein-Stancu operators and we call it as (p,q)-Bernstein-Stancu operators. We study approximation properties based on Korovkins type approximation theorem of (p,q)-Bernstein-Stancu operators. We also establish some direct theorems.

Keywords and phrases: (p,q)-integers; (p,q)-Bernstein operator; modulus of continuity; positive linear operator; Korovkin type approximation theorem.

AMS Subject Classifications (2010): 41A10, 41A25, 41A36, 40A30

1 Construction of Revised Operators

Mursaleen et. al [3] introduced (p,q)-analogue of Bernstein-Stancu operators as

$$S_{n,p,q}(f;x)=\sum_{k=0}^{n}\left[\begin{array}{c} n \\ k \end{array}\right]_{p,q} x^k \prod_{s=0}^{n-k-1}(p^s-q^s x)\ \ f\left(\frac{[k]_{p,q}+\alpha}{[n]_{p,q}+\beta}\right),\quad x\in[0,1]. \tag{1}$$

But $S_{n,p,q}(1;x)\neq 1$ for all $x\in[0,1)$ and $S_{n,p,q}(1;x)=1$ for $x=1$ only. Hence, we are re-introducing our operators as follows:

$$S_{n,p,q}(f;x)=\frac{1}{p^{\frac{n(n-1)}{2}}}\sum_{k=0}^{n}\left[\begin{array}{c} n \\ k \end{array}\right]_{p,q} p^{\frac{k(k-1)}{2}}x^k \prod_{s=0}^{n-k-1}(p^s-q^s x)\ \ f\left(\frac{p^{n-k}[k]_{p,q}+\alpha}{[n]_{p,q}+\beta}\right),\ \ x\in[0,1]. \tag{2}$$

Note that for $\alpha=\beta=0$, (p,q)-Bernstein-Stancu operators given by (2) reduces into (p,q)-Bernstein operators as given in [1, 2]. Also for $p=1$, (p,q)-Bernstein-Stancu operators given by (2) turn out to be q-Bernstein-Stancu operators.

For any $p>0$ and $q>0$, the (p,q) integers $[n]_{p,q}$ are defined by

$$[n]_{p,q}=p^{n-1}+p^{n-2}q+p^{n-3}q^2+...+pq^{n-2}+q^{n-1}=\begin{cases} \frac{p^n-q^n}{p-q}, & \text{when}\quad p\neq q\neq 1 \\ n\ p^{n-1}, & \text{when } p=q\neq 1 \\ [n]_q, & \text{when } p=1 \\ n, & \text{when } p=q=1 \end{cases}$$

where $[n]_q$ denotes the q-integers and $n = 0, 1, 2, \cdots$.

Obviously, it may be seen that $[n]_{p,q} = p^{n-1}[n]_{\frac{q}{p}}$.

We have the following basic result:

Lemma 1. For $x \in [0,1],\ 0 < q < p \leq 1$, we have

(i) $S_{n,p,q}(1;x) = 1$,
(ii) $S_{n,p,q}(t;x) = \frac{[n]_{p,q}}{[n]_{p,q}+\beta}x + \frac{\alpha}{[n]_{p,q}+\beta}$,
(iii) $S_{n,p,q}(t^2;x) = \frac{q[n]_{p,q}[n-1]_{p,q}}{([n]_{p,q}+\beta)^2}x^2 + \frac{[n]_{p,q}(2\alpha+p^{n-1})}{([n]_{p,q}+\beta)^2}x + \frac{\alpha^2}{([n]_{p,q}+\beta)^2}$.

Proof. (i)

$$S_{n,p,q}(1;x) = \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n} \left[\begin{array}{c} n \\ k \end{array} \right]_{p,q} p^{\frac{k(k-1)}{2}} x^k \prod_{s=0}^{n-k-1} (p^s - q^s x) = 1.$$

(ii)

$$\begin{aligned}
S_{n,p,q}(t;x) &= \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n} \left[\begin{array}{c} n \\ k \end{array} \right]_{p,q} p^{\frac{k(k-1)}{2}} x^k \prod_{s=0}^{n-k-1} (p^s - q^s x) \ \frac{p^{n-k}[k]_{p,q} + \alpha}{[n]_{p,q} + \beta} \\
&= \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n} \left[\begin{array}{c} n \\ k \end{array} \right]_{p,q} p^{\frac{k(k-1)}{2}} x^k \prod_{s=0}^{n-k-1} (p^s - q^s x) \ \frac{p^{n-k}[k]_{p,q}}{[n]_{p,q} + \beta} \\
&+ \frac{\alpha}{[n]_{p,q} + \beta} \frac{1}{p^{\frac{n(n-1)}{2}}} \sum_{k=0}^{n} \left[\begin{array}{c} n \\ k \end{array} \right]_{p,q} p^{\frac{k(k-1)}{2}} x^k \prod_{s=0}^{n-k-1} (p^s - q^s x) \\
&= \frac{[n]_{p,q}}{p^{\frac{n(n-3)}{2}}} \sum_{k=0}^{n-1} \left[\begin{array}{c} n-1 \\ k \end{array} \right]_{p,q} p^{\frac{(k+1)(k)}{2}} x^{k+1} \prod_{s=0}^{n-k-2} (p^s - q^s x) \frac{1}{p^{k+1}([n]_{p,q} + \beta)} + \frac{\alpha}{[n]_{p,q} + \beta} \\
&= \frac{[n]_{p,q}}{[n]_{p,q} + \beta} x \ \frac{1}{p^{\frac{(n-1)(n-2)}{2}}} \sum_{k=0}^{n-1} \left[\begin{array}{c} n-1 \\ k \end{array} \right]_{p,q} p^{\frac{k(k-1)}{2}} x^k \prod_{s=0}^{n-k-2} (p^s - q^s x) + \frac{\alpha}{[n]_{p,q} + \beta} \\
&= \frac{[n]_{p,q}}{[n]_{p,q} + \beta} x + \frac{\alpha}{[n]_{p,q} + \beta}.
\end{aligned}$$

(iii)

$$\begin{aligned} S_{n,p,q}(t^2;x) &= \frac{1}{p^{\frac{n(n-1)}{2}}}\sum_{k=0}^{n}\left[\begin{array}{c} n \\ k \end{array}\right]_{p,q} p^{\frac{k(k-1)}{2}}x^k\prod_{s=0}^{n-k-1}(p^s-q^sx)\ \left(\frac{p^{n-k}[k]_{p,q}+\alpha}{[n]_{p,q}+\beta}\right)^2 \\ &= \frac{1}{([n]_{p,q}+\beta)^2}\ \frac{1}{p^{\frac{n(n-1)}{2}}}\Bigg[p^{2n}\sum_{k=0}^{n}\left[\begin{array}{c} n \\ k \end{array}\right]_{p,q} p^{\frac{k(k-1)}{2}}x^k\prod_{s=0}^{n-k-1}(p^s-q^sx)\ \frac{[k]_{p,q}^2}{p^{2k}} \\ &\quad +2\alpha\ p^n\sum_{k=0}^{n}\left[\begin{array}{c} n \\ k \end{array}\right]_{p,q} p^{\frac{k(k-1)}{2}}x^k\prod_{s=0}^{n-k-1}(p^s-q^sx)\ \frac{[k]_{p,q}}{p^k} \\ &\quad +\alpha^2\ \sum_{k=0}^{n}\left[\begin{array}{c} n \\ k \end{array}\right]_{p,q} p^{\frac{k(k-1)}{2}}x^k\prod_{s=0}^{n-k-1}(p^s-q^sx)\Bigg]. \end{aligned}$$

On shifting the limits and using $[k+1]_{p,q}=p^k+q[k]_{p,q}$, we get our desired result.

Lemma 2.2. For $x\in[0,1],\ 0<q<p\leq 1$

(i) $S_{n,p,q}\big((t-x);x\big)=\frac{\alpha-\beta x}{[n]_{p,q}+\beta}$,
(ii) $S_{n,p,q}\big((t-x)^2;x\big)=\Big\{\frac{q[n]_{p,q}[n-1]_{p,q}-[n]_{p,q}^2+\beta^2}{([n]_{p,q}+\beta)^2}\Big\}x^2+\Big\{\frac{p^{n-1}[n]_{p,q}-2\alpha\beta}{([n]_{p,q}+\beta)^2}\Big\}x+\frac{\alpha^2}{([n]_{p,q}+\beta)^2}$.

Proof. (ii) By linearity of the operator we have

$$\begin{aligned} S_{n,p,q}\big((t-x)^2;x\big) &= S_{n,p,q}(t^2;x)-2xS_{n,p,q}(t;x)+x^2S_{n,p,q}(1;x) \\ &= \frac{q[n]_{p,q}[n-1]_{p,q}}{([n]_{p,q}+\beta)^2}x^2+\frac{[n]_{p,q}(2\alpha+p^{n-1})}{([n]_{p,q}+\beta)^2}x \\ &\quad +\frac{\alpha^2}{([n]_{p,q}+\beta)^2}-2x\frac{[n]_{p,q}}{[n]_{p,q}+\beta}-2x\frac{\alpha}{[n]_{p,q}+\beta}+x^2 \\ &= \left\{\frac{q[n]_{p,q}[n-1]_{p,q}-[n]_{p,q}^2+\beta^2}{([n]_{p,q}+\beta)^2}\right\}x^2+\left\{\frac{p^{n-1}[n]_{p,q}-2\alpha\beta}{([n]_{p,q}+\beta)^2}\right\}x+\frac{\alpha^2}{([n]_{p,q}+\beta)^2}. \end{aligned}$$

2 Main Results:

2.1 Korovkin type approximation theorem

Remark 2.1 *For $q\in(0,1)$ and $p\in(q,1]$, it is obvious that $\lim_{n\to\infty}[n]_{p,q}=0$ or $\frac{1}{p-q}$. In order to reach to convergence results of the operator $L_{p,q}^n(f;x)$, we take a sequence $q_n\in(0,1)$ and $p_n\in(q_n,1]$ such that $\lim_{n\to\infty}p_n=1$, $\lim_{n\to\infty}q_n=1$ and $\lim_{n\to\infty}p_n^n=1$, $\lim_{n\to\infty}q_n^n=1$. So we get $\lim_{n\to\infty}[n]_{p_n,q_n}=\infty$.*

Theorem 3.1.1. Let sequence q_n and p_n be such that it satisfies Remarks 2.1, then $S_{n,p,q}(f,x)$ converges uniformly to f on $[0,1]$.

Proof. By the Bohman-Korovkin Theorem it is sufficient to show that

$$\lim_{n\to\infty} \|S_{n,p_n,q_n}(t^m;x) - x^m\|_{C[0,1]} = 0, \quad m = 0, 1, 2.$$

By Lemma 2.1 (i)-(ii), it is clear that

$$\lim_{n\to\infty} \|S_{n,p_n,q_n}(1;x) - 1\|_{C[0,1]} = 0;$$

$$\lim_{n\to\infty} \|S_{n,p_n,q_n}(t;x) - x\|_{C[0,1]} = 0.$$

Using $q_n[n-1]_{p_n,q_n} = [n]_{p_n,q_n} - p_n^{n-1}$ and by Lemma 2.1(iii), we have

$$|S_{n,p_n,q_n}(t^2;x) - x^2|_{C[0,1]} = \left| \frac{q[n]_{p,q}[n-1]_{p,q}}{([n]_{p,q}+\beta)^2}x^2 + \frac{[n]_{p,q}(2\alpha + p^{n-1})}{([n]_{p,q}+\beta)^2}x + \frac{\alpha^2}{([n]_{p,q}+\beta)^2} - x^2 \right|.$$

Taking maximum of both sides of the above inequality, we get

$$\|S_{n,p_n,q_n}(t^2;x) - x^2\|_{C[0,1]} \leq \frac{q[n]_{p,q}[n-1]_{p,q}}{([n]_{p,q}+\beta)^2} - 1 + \frac{[n]_{p,q}(2\alpha + p^{n-1})}{([n]_{p,q}+\beta)^2} + \frac{\alpha^2}{([n]_{p,q}+\beta)^2}$$

which yields

$$\lim_{n\to\infty} \|S_{n,p_n,q_n}(t^2;x) - x^2\|_{C[0,1]} = 0.$$

Thus the proof is completed and all other results follow similarly.

References

[1] M. Mursaleen, Khursheed J. Ansari, Asif Khan, On (p, q)-analogue of Bernstein operators, Applied Mathematics and Computation, 266 (2015) 874-882.

[2] M. Mursaleen, Khursheed J. Ansari, Asif Khan, On (p, q)-analogue of Bernstein operators (revised), *arxiv*:1503.07404v2[math CA].

[3] M. Mursaleen, K. J. Ansari and Asif Khan,, Some Approximation Results by (p, q)-analogue of Bernstein-Stancu Operators, *Applied Mathematics and Computation* 264,(2015), 392-402.